

Phyllanthus niruri Linn increase infiltrating lymphocyte and apoptosis of rectosigmoid cancer patients: A Phase II Clinical Trial

by Ignatius Riwanto

Submission date: 31-Jan-2019 12:09PM (UTC+0700)

Submission ID: 1070938658

File name: Artikel_C2_Phyllanthus_niruri_Linn_increase....pdf (260.43K)

Word count: 4148

Character count: 22796

13

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/319983095>

Phyllanthus niruri Linn increase infiltrating lymphocyte and apoptosis of rectosigmoid cancer patients. A phase II clinical trial

Article · July 2017

DOI: 10.15562/bmj.v6i3.674

CITATIONS

0

READS

23

7 authors, including:



Toar Mambu
Dr. D. Kandou General Hospital, Manado, Indonesia

2 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



DiscoverSys

Published by DiscoverSys

Phyllanthus niruri Linn increase infiltrating lymphocyte and apoptosis of rectosigmoid cancer patients: A Phase II Clinical Trial



CrossMark

Ignatius Riwanto,^{1*} Parish Budiono,¹ Abdul Mughni,¹ Martahadinan,²
Nugrahanta Dasa Putra,² Toar Deliezer Bram Mambu,² Mochamad Sayuti²

ABSTRACT

Background: Phyllanthus niruri Linn (PNL) extract has potential cytotoxic effects on colorectal cancer in both in vitro and in vivo studies, but this has never been reported in human subjects. The purpose of this study is to elaborate whether PNL extract also has cytotoxic effects on human colorectal cancer.

Method: A one-group pre- and post-test analysis was conducted on 15 patients with non-obstructive operable rectosigmoid cancer without neo-adjuvant chemotherapy or chemoradiation. Before commencing treatment using PNL extract (100 mg Stimuno), all patients underwent rectoscopy biopsy. Stimuno treatment was given for at least 14 days and later patients underwent rectal resection. Specimens from biopsy and rectosigmoid resection were managed by the Department of Anatomical Pathology, Faculty of Medicine,

Gajahmada University, Yogyakarta, for evaluating the diagnosis, infiltrating lymphocyte, apoptosis index, perforin, and granzyme-B and caspase-3 expression. A paired *t*-test was done to evaluate the significant increase in measured variables, and a linear regression analysis was also done to evaluate the role of individual variables.

Result: Infiltrating lymphocyte; expression of perforin, granzyme-B, and caspase-3; and apoptosis index significantly ($p = 0.000$) improved after the treatment using PNL extract. After conducting the linear regression analysis, we found that the only variable that significantly correlated with apoptosis was infiltrating lymphocyte.

Conclusion: Use of PNL extract increased the apoptosis index of colorectal cancer through an increase in infiltrating lymphocyte.

Keyword: Phyllanthus niruri Linn, rectal cancer, infiltrating lymphocyte, apoptosis

Cite This Article: Riwanto, I., Budiono, P., Mughni, A., Martahadinan, Putra, N.D., Mambu, T.D.B., Sayuti, M. 2017. Phyllanthus niruri Linn increase infiltrating lymphocyte and apoptosis of rectosigmoid cancer patients: A Phase II Clinical Trial. *Bali Medical Journal* 6(3): 509-513. DOI: [10.15562/bmj.v6i3.674](https://doi.org/10.15562/bmj.v6i3.674)

¹Staff Dept. of Surgery, Digestive Division, Dr. Kariadi Hospital, Faculty of Medicine, University of Diponegoro

²Student of Digestive Surgery Subspecialist Program

INTRODUCTION

Neo-adjuvant or adjuvant treatment of colorectal cancer has reduced the recurrence rate and improved the 5-year survival rate. A study of 119,363 colon cancer cases between January 1991 and December 2000 showed that according to American Joint Committee on Cancer (AJCC) TNM System Sixth Edition stages, the 5-year-stage-specific survival rate was 93.2% for stage I, 84.7% for stage IIa, 72.2% for stage IIb, 83.4% for stage IIIa, 64.1% for stage IIb, 44.3% for stage IIc, and 8.1% for stage IV. The significantly better result of stage IIIa compared to that of Stage II could be attributed to the use of chemotherapy. National Institutes of Health (NIH) 1990 Consensus Conference guidelines for colorectal cancer recommend adjuvant chemotherapy for stage III colon cancer but not for stage II patients.¹ Review done by El Zouhairi et al. 2011² showed that over the past decade, the survival rate of metastatic colorectal cancer patients³ proved dramatically after using chemotherapy. In the mid-1990s, the median overall survival rate of those treated with 5-FU-based regimen was about 12 months; the combination of 5-FU with irinotecan or oxaliplatin

increased the OS to approximately 18%, and after addition with targeted therapy, the OS approached 30% in some studies. Although there was a lot of improvement in survival, the results of stage IIb, IIc, and IV were still disappointing.

Thus, these disappointing results that came about even after the treatment using modern chemo for targeted therapy of metastatic colorectal cancer stimulated research into finding substances that may solve this problem. Immunomodulator seems to be the other alternative to improve survival and reduce the recurrence of colorectal cancer. Phyllanthus niruri Linn (PNL) is one of the potent immunomodulator; its efficacy has already been proven in relation to its capacity to inhibit the development of experimental colorectal cancer of Sprague-Dawley male rat. PNL extract significantly decreased AgNORs, the cancer cells' proliferative marker, and macrophagic tumor growth.³ In vitro study showed that spray-dried extracts of Phyllanthus niruri (SDEPN) has a significantly better cytotoxic effect on Hepatocellular cell cancer line (Hep G2) but not on colorectal cancer cell line

*Correspondence to :
Ignatius Riwanto
iriwanto@gmail.com
Mobile phone: +62811277088

Received: 2017-06-16
Accepted: 2017-08-28
Published: 2017-09-1

Open access: www.balimedicaljournal.org and ojs.unud.ac.id/index.php/bmj

(HT29) compared to control. However, pretreatment with SDEPN followed by cisplatin had a synergistic effect and showed a significantly much better cytotoxic effect on HT 29.⁴ PNL extract has already been used widely as traditional medicine in Indonesia. In experimental animal model PNL is used for the treatment of kidney or liver, even in extremely high doses at >5000 mg/kg. b.w.⁵

The purpose of this study is to evaluate whether PNL extract can also influence the growth of colorectal cancer patients through perforin-granzyme pathway, by measuring infiltrating lymphocyte, perforin, granzyme-B, caspase-3, and apoptosis index before and after the PNL extract treatment.

METHOD

The study sample included patients with resectable non-obstructive rectal cancer but who did not undergo neo-adjuvant chemotherapy or chemoradiation; other selection criteria included the following: age above 40 years, a BMI score that is more than 20, and hemoglobin content that is more than 10 gr%. Patients were explained about the study, and informed consent was obtained from them; eventually, a total of 15 rectal cancer patients were enrolled in the present study which was conducted in Dr. Kariadi Hospital, Semarang Indonesia, between May and July 2016. The study has already been approved by the Ethical Clearance Committee of Dr. Kariadi Hospital, Faculty of Medicine, Diponegoro University. The study was based on pre-test and post-test design. All patients underwent rectoscopy and biopsy and were managed as outpatients. The biopsy specimen was fixed with formalin buffer within 18–24 hours and further processed with paraffin block. The paraffin block was sent to Anatomical Pathology Laboratory of Faculty of Medicine, Gajahmada University, for further analysis regarding the microscopic diagnosis and immune histochemical staining for detecting perforin, granzyme-B, caspase-3 expression, and degree of apoptosis of cancer cells. The next day after rectoscopy the patient was treated with PNL extract (R/Stimuno) 100 mg/day for a minimum 14 days. Stimuno contains a single composition formula of *Phyllanthus niruri* plant extracts taken from roots, leaves, stems, and fruit and formulated as capsules. Stimuno is being used since 1999; we obtained a certificate of phytopharmaca by POM RI on March 22, 2005. After Stimuno treatment for 14 days or more, patients were admitted to Dr. Kariadi Hospital and underwent rectal resection, and specimens were treated as biopsy specimen in the same Anatomical Pathology Laboratory. Infiltrating lymphocyte was measured on Hematoxylin Eosin

staining specimen under 400x power field. The apoptosis index was measured using the terminal deoxynucleotide transferase dUTP nick-end labeling method. Perforin, granzyme-B, and caspase-3 expressions were measured on immune histochemical staining specimen under 400x power field. All were measured by two independent anatomical pathology experts using the same criteria and clinical agreement was more than 95%. As the numerical data showed normal distribution, paired *t*-test was used to compare variables used in the study before and after treatment. Linear regression analysis of the increasing value of variables was done to identify which variable significantly predicted the increased apoptosis.

RESULT

All the 15 rectal or sigmoid cancer patients completed the study. The mean length of treatment with stimuno was 17 days (14–22 days). Patients gender classification was as follows: 9 males and 6 females; mean of age of the patients was 53.66 ± 14.52 years. Mean serum urea level pre-treatment was 27.8 ± 6.48 mg/dl, range 17–38, and post-treatment the same was 29.9 ± 7.78 mg/dl, range 19–42 ($p = 0.000$). Mean serum creatinine pre-treatment was 0.89 ± 0.22 mg/dl, range 0.56–1.20, and post-treatment it was 0.85 ± 0.18 mg/dl, range 0.56–1.35 ($p = 0.000$). Serum alanine aminotransferase (ALT) pre-treatment was 28.3 ± 7.71 unit/l, range 17–36, and post-treatment it was 26.8 ± 6.38 unit/l, range 17–40 ($p = 0.000$).

Infiltrating Lymphocyte, granzyme-B, perforin, caspase-3 expression, and apoptosis index significantly increased after treatment with PNL extract (Table 1).

To know the relationship among variables, since it was a pre- and post-test design, a correlation matrix of increasing value of variables (the

Table 1 Variables before and after PNL extract treatment *paired *t*-test

No	Variables		Mean (SD)	p*
1	Lymphocyte	pre	45.4 (4.8)	0.000
		post	179.8 (38.8)	
2	Granzyme-B	pre	25.4 (4.8)	0.000
		post	65.7 (7.9)	
3	Perforin	pre	37.6 (20.7)	0.000
		post	42.0 (20.9)	
4	Caspase-3	pre	37.4 (13.2)	0.000
		post	61.0 (18.3)	
5	Apoptosis	pre	16.9 (5.0)	0.000
		post	87.2 (20.3)	

Table 2 Correlation matrix among increasing variables. *r*: Pearson Correlation, *p* (significant level two-tail), *Infiltrating

Variables		*Lymphocyte	Granzyme-B	Perforin	Caspase-3	Apoptosis
*Lymphocyte	<i>r</i>	1	0.15	-0.03	0.46	0.71
	<i>p</i>	0.589	0.912	0.080	0.003	
Granzyme-B	<i>r</i>	0.15	1	0.25	-0.12	0.18
	<i>p</i>	0.589	0.363	0.667	0.502	
Perforin	<i>r</i>	-0.03	0.25	1	-0.08	0.14
	<i>p</i>	0.912	0.363	0.753	0.613	
Caspase-3	<i>r</i>	0.46	-0.12	-0.08	1	0.31
	<i>p</i>	0.08	0.667	0.753	0.255	
Apoptosis	<i>r</i>	0.71	0.18	0.14	0.31	1
	<i>p</i>	0.003	0.502	0.613	0.255	

Table 3 Linear regression with dependent variables increasing apoptosis, while increasing infiltrating lymphocyte, granzyme-B, perforin, and caspase-3 expression were independent variables

Model Significance	Unstandardized Coefficients	Standardized Coefficients	<i>t</i>	
	B	SE	Beta	
Constant	17.753	14.845	1.196	0.253
Lymphocyte diff.	0.391	0.106	0.715	3.690

different between post- and pre-test values) was done. It showed that the one variable that strongly correlated with apoptosis was lymphocyte ($r = 0.715$, $p = 0.003$). A nearly significant correlation was shown between infiltrating lymphocyte and caspase-3 ($r = 0.46$, $p = 0.080$; Table 2). On linear regression, it was more clear that apoptosis was strongly correlated with only lymphocyte (Table 3).

DISCUSSION

This study was phase II clinical trial, and the study purpose was to know whether the extract of PNL, which was already proven to be effective in arresting the growth of experimental Windstar colorectal cancer,⁵ also has a positive effect on arresting human colorectal cancer growth. The research regarding PNL's capacity to arrest the growth of human colorectal cancer cells has never been reported. Since PNL extract has already been used widely in the Indonesian community as a traditional medicine and proven safe, the Phase I clinical trial was not done. This argument is supported by the result of this study. Although serum of urea, creatinine, and ALT changed significantly, the levels of both pre- and post-PNL treatment were still within normal range. It means that PNL is safe for kidney and liver, which is in accordance with the previous research⁵. The results of the study showed that PNL extract significantly increased the infiltrating lymphocyte, granzyme-B, perforin, caspase-3

expression, and cancer cell apoptosis. These results are supported by several studies that used PNL in arresting the development of various cancer cells.

Past studies using PNL on hepatic carcinoma or prostatic cancer cell line reported results that agree with the findings of this study. Cell cytotoxicity assays of *Phyllanthus niruri* extract showed that the active component is corilagin, which has a broad-spectrum antitumor activity to arrest hepatic carcinoma cell line, better antitumor potential, and lower toxicity in normal cells.⁶ The other study on hepatic carcinoma cell line showed that SDEPN reduces the viable liver cancer cell and increases apoptosis.⁴ Both aqueous and methanolic extracts of PNL significantly increase pro-apoptotic protein (Bax) and decrease anti-apoptotic protein (Bcl-2) of the prostate cancer (PC-3) cell line. NfKB also decreases significantly with both aqueous and methanolic extract of PNL. Bax protein could induce cytochrome c release from mitochondria, which can then induce proteolytic activation of procaspase-9. This in turn activates caspase-3 and finally leads to induction of apoptosis in PC-3 cells.⁷

Studies that evaluated the effect of PNL on colorectal cancer have used only animal experimental model and have been conducted only in vitro. Study on Wistar rat showed that use of 1,2 DMH (Dimethylhydrazine) led to the development of colorectal cancer and that PNL extracts reduced the AgNOR as a marker of cell proliferative and also

reduced the size of the tumor.³ In combination with cisplatin, SDEPN had a synergistic effect and showed significantly much better cytotoxic effect on colorectal cancer cell line (HT 29) compared to cisplatin or SDEPN alone.⁸ The other study for colorectal cancer²¹ was using *Phyllanthus watsonii*. In vitro study on human¹⁰ gynecologic and colon cancer cells line showed that *P. watsonii* arrested cell cycle at different²⁹ growth phases and increased the Caspase-3 activity as a single agent or in¹⁰ combination with doxorubicin. It can be concluded that the sub-fraction of *P. watsonii* strongly inhibits the growth and induces apoptosis²¹ of human gynecologic and colorectal cancer.⁹ Downregulation of anti-apoptotic protein (BCL-2) is comparable between *P. watsonii* and *P. niruri* on the study⁸ done in vitro with lung cancer cell line.¹⁰

There are two main apoptotic pathways: the extrinsic or death receptor pathway and the intrinsic or mitochondrial pathway. There is an additional pathway that involves T-cell-mediated cytotoxicity and perforin-granzyme-dependent killing of the cell.¹¹ In this study, the role of PNL extract as an apoptotic agent was measured through perforin-granzyme-dependent killing pathway. The strong correlation was only shown between infiltrating lymphocyte and apoptosis index. The relationship between infiltrating lymphocyte and caspase-3 was almost significant; meanwhile, increased expression of perforin and granzyme-B was not significantly correlated to the apoptosis and also to infiltrating lymphocyte. Granzymes that play an important role in the apoptosis mechanism are granzyme A and granzyme B. The extrinsic, intrinsic, and granzyme B pathways converge on the same execution pathway that is the cleavage of caspase-3, as executor of apoptosis. Caspase-6 and -7 are the other executors of apoptosis. The granzyme A pathway activates a parallel, caspase-independent cell death pathway via single-stranded DNA¹⁴ damage. Granzyme A cannot activate caspase but can kill cancer cells by the direct cleavage of nuclear proteins, thereby facilitating the formation and accumulation of single-stranded DNA breaks.¹¹ In this study, granzyme A was not measured, and measuring infiltrating lymphocyte as a whole means it consists of many types of subset lymphocyte; meanwhile, lymphocytes²² are capable of initiating apoptosis are only cytotoxic T lymphocytes (CTLs) and natural killer (NK) cells. To kill the cell target needs direct contact. Therefore, only a part of infiltrating lymphocyte expresses perforin and granzyme.¹² The above explanation makes it evident that an increase in PNL content arrests the increase in infiltrating lymphocyte and apoptosis of rectosigmoid cancer partly through perforin and granzyme-B pathways and granzyme-A, but extrinsic pathway still needs to be studied further. Further studies regarding the

role of PNL needs to be more specific to lymphocyte⁴ CTLs, and NK cells should be carried out.

The extrinsic signaling pathways that initiate apoptosis involve transmembrane receptor-mediated interactions. These involve death receptors that are members of the tumor necrosis factor (TNF) receptor gene superfamily.¹¹ Study³⁴ done by Reissfelder et al. (2015)¹³ showed that upregulation of TNF- α expression in tumor-infiltrating lymphocytes (TILs) strongly correlated with an increase in the total amount of intra-tumoral TNF- α , which is indicative of tumor-specific CTL activity. Increased TNF- α concentration was an independent prognostic factor. It suggests that intra-tumoral TNF- α might be a more suitable factor for prognosis prediction in patients with CRC than the quantification of effector TC infiltrates.

The relationship between infiltrating lymphocyte and the survival of colorectal cancer has already been studied. The low tumor-infiltrating lymphocyte (TIL) group was significantly correlated with a poorly differentiated status and perineural invasion. During the median 54-month follow-up period, the low-TIL group had a significantly lower five-year overall survival and disease-free survival rates than the high-TIL group of patients with stage III colorectal cancer ($p = 0.005$ and $p = 0.03$, respectively).¹⁴ Meta-analysis of 12 publications regarding tumor inflammatory infiltrate showed that high generalized tumor inflammatory infiltrate was associated with good overall survival (HR: 0.59, 95% CI: 0.48–0.72), cancer-specific survival (HR: 0.40, 95% CI: 0.27–0.61), and disease-free survival (HR: 0.72, 95% CI: 0.57–0.91).¹⁵

The circulating lymphocyte¹ also has a role as a prognostic factor. The low pre-treatment lymphocyte-to-monocyte Ratio (LMR) group had a significantly worse overall survival rate ($p = 0.0011$). Moreover, patients who demonstrated low pre-treatment LMR and normalization after treatment exhibited a better overall survival rate than the patients with low pre- and post-treatment LMR values. Lymphocytes play an important role in the antitumor immunity of the host, including cytotoxic cell death and the inhibition¹ of tumor cell proliferation and migration. On the other hand, monocytes play an important role in tumor progression and metastasis. Tumor-associated macrophages (TAMs), which are derived from circulating monocytes, suppress adaptive immunity and promote angiogenesis, invasion, migration, and tumor growth.¹⁶ Significantly worse disease-free survival can be observed in patients with lower LMR levels (<3.78) using uni- and multi-variate analyses ($p = 0.01$ and $p = 0.015$, respectively).¹⁷ The relationship between circulating lymphocyte and infiltrating lymphocyte in colorectal cancer is reported by Pane et al. (2015).¹⁸ Neutrophil-to-lymphocyte ratio (NLR) of

19 more results in significantly worse prognosis, higher pT- and pN-stage, and a greater incidence of extramural venous invasion compared to NLR below 5; meanwhile, moderate-to-severe presence of infiltrating lymphocytes in tumor margin is a significantly better prognosis compared to total absence or a mild presence. Infiltrating lymphocyte is also significantly higher in patients with NLR below 5. In this study, the circulating lymphocyte was not measured, and its impact on survival was not evaluated thus. The PNL was increasing the infiltrating of the rectosigmoid cancer; therefore, on the basis of Pane's study results,¹⁸ we assume that PNL will also increase the circulating lymphocyte and could enhance the survival rate. Further study is needed to elaborate our hypotheses.

Since our study had only a one-group pre- and post-test design, the history and maturity factor may affect the results of the study. This is the main limitation of this study. Therefore, the next study should be randomized control trial (Phase III clinical trial) that makes maturity and history factors comparable and individual²⁰ effect of the PNL extract can be measured. However, results of this study support²⁰ potential role of PNL as an immune modulator in the treatment of rectosigmoid cancer, but further research should be carried out to study in depth the role of PNL extract in arresting the cancer cell growth. In conclusion, it is clear that PNL extract improves the infiltrating lymphocyte and increases the apoptosis index, partly through perforin-granzyme-B pathway in patients with rectosigmoid cancer.

ACKNOWLEDGMENT

17 express our sincere gratitude to the Head of the Department of Anatomical Pathology, Faculty of Medicine, University of Gajahmada and his staff, for managing the rectal specimen and measuring the studied variables. Also we express²⁵ our gratitude to the Digestive Division, Dept. of Surgery, Dr. Kariadi Hospital, Faculty of Medicine, Diponegoro University, for funding the research.

REFERENCES

- O'Connell JB, Maggard MA, Ko CY. Colon Cancer Survival Rates With the New American Joint Committee on Cancer Sixth Edition Staging. *J Natl Cancer Inst* 2004;96(19):1420-5. DOI: [10.1093/jnci/djh275](https://doi.org/10.1093/jnci/djh275)
- El Zouhairi M, Charabaty A, Pishvaian. Molecularly Targeted Therapy for Metastatic Colon Cancer: Proven Treatments and Promising New Agents. *Gastrointest Cancer Res* 2011;4:15-21
- Savitri E, Riwanto I, Tjahjono, Dharmana E. Ekstrak Phyllanthus Niruri Linn. Pertumbuhan tumor dan proliferasi sel kanker kolorektal: Studi eksperimental pada tikus Sprage-Dawley yang diinduksi 1,2 DMH. *M. Med. Indones*. 2012;46(1):25-32.
- Junior RFA, Soares LAL, da Costa Porto CR, de Aquino RGF, Guedes HG, Petrovick PR et al. Growth inhibitory effects of Phyllanthus niruri extracts in combination with cisplatin on cancer cell lines. *World J Gastroenterol* 2012 August 21; 18(31): 4162-4168. DOI: [10.3748/wjg.v18.i31.4162](https://doi.org/10.3748/wjg.v18.i31.4162)
- Asare GA, Addo P, Bgyei K, Gyan B, Adjei S, Out-Nyarko LS et al. Acute toxicity studies of aqueous leaf extract of Phyllanthus niruri. *Interdiscip Toxicol*. 2011; Vol. 4(4): 206-210. DOI: [10.2478/v10102-011-0031-9](https://doi.org/10.2478/v10102-011-0031-9)
- Zheng ZZ, Chen LH, Liu SS, Deng Y, Zheng GH, Gu Y et al. Bioguided Fraction and Isolation of the Antitumor Components from Phyllanthus niruri L. *BioMed Research International* 2016; Article ID 9729275: 1-7 <http://dx.doi.org/10.1155/2016/9729275>
- Tang YQ, Jaganath IB, Manikam R, Sekaran SD. Phyllanthus Suppresses Prostate Cancer Cell, PC-3, Proliferation and Induces Apoptosis through Multiple Signalling Pathways (MAPKs, PI3K/Akt, NF- κ B, and Hypoxia). *Evidence-Based Complementary and Alternative Medicine* 2013; Article ID 609581: 1-13 <http://dx.doi.org/10.1155/2013/609581>
- Junior RFA, de Souza TP, Pires JGL, Soares LAL, de Araujo AA, Petrovick PR et al. A dry extract of Phyllanthus niruri protects normal cells and induces apoptosis in human liver carcinoma cells *Experimental Biology and Medicine* 2012; 237: 1281-1288. DOI: [10.1258/ebm.2012.012130](https://doi.org/10.1258/ebm.2012.012130)
- Ramasamy S, Abdul Wahab N, Zainal Abidin N, Manickam S, Zakaria Z. Growth Inhibition of Human Gynecologic and Colon Cancer Cells by Phyllanthus watsonii through Apoptosis Induction. *PLoS ONE* 2012;7(4): e34793. DOI: [10.1371/journal.pone.0034793](https://doi.org/10.1371/journal.pone.0034793)
- Lee SH, Jaganath IB, Manikam R, Sekaran SD. Inhibition of Raf-MEK-ERK and Hypoxia pathways by Phyllanthus prevents metastasis in human lung (A549) cancer cell line. *Complementary and Alternative Medicine* 2013, 13:271. <http://www.biomedcentral.com/1472-6882/13/271>
- Elmore S. Apoptosis: A Review of Programmed Cell Death. *Toxicol Pathol*. 2007; 35(4): 495-516
- Trapani JA, Smyth MJ. Functional Significance Of The Perforin/Granzyme Cell Death Pathway. *Immunology* 2002;2: 735-747. DOI: [10.1038/nri911](https://doi.org/10.1038/nri911)
- Reissfelder C, Stamova S, Gossmann C, Braun M, Bonertz A, Walliczek U et al. Tumor-specific cytotoxic T lymphocyte activity determines colorectal cancer patient prognosis. *J Clin Invest*. 2015;125(2):739-751. DOI: [10.1172/JCI74894](https://doi.org/10.1172/JCI74894)
- Huh JW, Lee JH, Kim HR. Prognostic Significance of Tumor-Infiltrating Lymphocytes for Patients With Colorectal Cancer. *Arch Surg*. 2012;147(4):366-371. <http://jamanetwork.com/pdfaccess.ashx?url=/data/journals/surg/23388/ on 02/09/2017>
- Mei Z, Liu Y, Liu C, Cui A, Liang Z, Wang G et al. Tumour-infiltrating inflammation and prognosis in colorectal cancer: systematic review and meta-analysis. *British Journal of Cancer* (2014) 110, 1595-1605. DOI: [10.1038/bjc.2014.46](https://doi.org/10.1038/bjc.2014.46)
- Shibutani M, Maeda K, Nagahara H, Ohtani H, Sakurai K, Yamazoe S, et al. Prognostic significance of the lymphocyte-to-monocyte ratio in patients with metastatic colorectal cancer. *World J Gastroenterol* 2015 September 14; 21(34): 9966-9973. DOI: [10.3748/wjg.v21.i34.9966](https://doi.org/10.3748/wjg.v21.i34.9966)
- Xiao WW, Zhang LN, You KY, Huang R, Yu X, Ding PR et al. A Low Lymphocyte-to-Monocyte Ratio Predicts Unfavorable Prognosis in Pathological T3N0 Rectal Cancer Patients Following Total Mesorectal Excision. *Journal of Cancer* 2015; 6(7): 616-622. DOI: [10.7150/jca.117](https://doi.org/10.7150/jca.117)
- Pine JK, Morris E, Hutchins GG, West NP, Jayne DG, Quirke et al. Systemic neutrophil-to-lymphocyte ratio in colorectal cancer: the relationship to patient survival, tumour biology and local lymphocytic response to tumour *British Journal of Cancer* (2015) 113, 204-211. DOI: [10.1038/bjc.2015.87](https://doi.org/10.1038/bjc.2015.87)



This work is licensed under a Creative Commons Attribution

Phyllanthus niruri Linn increase infiltrating lymphocyte and apoptosis of rectosigmoid cancer patients: A Phase II Clinical Trial

ORIGINALITY REPORT

25%

SIMILARITY INDEX

20%

INTERNET SOURCES

20%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1

office.wjgnet.com

Internet Source

3%

2

Submitted to iGroup

Student Paper

2%

3

jnci.oxfordjournals.org

Internet Source

2%

4

www.ncbi.nlm.nih.gov

Internet Source

2%

5

jsurg.com

Internet Source

1%

6

6jgc.com.www.mobile.jci.org

Internet Source

1%

7

mdanderson.influuent.utsystem.edu

Internet Source

1%

8

krishikosh.egranth.ac.in

Internet Source

1%

9	Mei, Z, Y Liu, C Liu, A Cui, Z Liang, G Wang, H Peng, L Cui, and C Li. "Tumour-infiltrating inflammation and prognosis in colorectal cancer: systematic review and meta-analysis", British Journal of Cancer, 2014. Publication	1%
10	journals.plos.org Internet Source	1%
11	www.scribd.com Internet Source	1%
12	Tang, Yin-Quan, InduBala Jaganath, Rishya Manikam, and Shamala Devi Sekaran. "Phyllanthus Suppresses Prostate Cancer Cell, PC-3, Proliferation and Induces Apoptosis through Multiple Signalling Pathways (MAPKs, PI3K/Akt, NFB, and Hypoxia)", Evidence-based Complementary and Alternative Medicine, 2013. Publication	1%
13	doc.rero.ch Internet Source	1%
14	www.columbia.edu Internet Source	1%
15	www.spandidos-publications.com Internet Source	1%

16

Submitted to Symbiosis International University

Student Paper

<1 %

17

edoc.ub.uni-muenchen.de

Internet Source

<1 %

18

www.internationalsurgery.org

Internet Source

<1 %

19

eprints.whiterose.ac.uk

Internet Source

<1 %

20

Oncology, 2006.

Publication

<1 %

21

"Anticancer Plants: Mechanisms and Molecular Interactions", Springer Nature, 2018

Publication

<1 %

22

preview-bmcimmunol.biomedcentral.com

Internet Source

<1 %

23

Farida, Helmia, Juliëtte A. Severin, M. Hussein Gasem, Monique Keuter, Hendro Wahyono, Peterhans van den Broek, Peter W. M. Hermans, and Henri A. Verbrugh.

"Nasopharyngeal Carriage of Streptococcus pneumonia in Pneumonia-Prone Age Groups in Semarang, Java Island, Indonesia", PLoS ONE, 2014.

Publication

<1 %

24

onlinelibrary.wiley.com

<1 %

25

ejournal2.undip.ac.id

Internet Source

<1 %

26

doaj.org

Internet Source

<1 %

27

Submitted to National postgraduate Medical
College of Nigeria

Student Paper

<1 %

28

iem.modares.ac.ir

Internet Source

<1 %

29

Tan, Li, Zhang Zhang, Donglin Gao, Jinfeng
Luo, Zheng-Chao Tu, Zhengqiu Li, Lijie Peng,
Xiaomei Ren, and Ke Ding. "4-Oxo-1, 4-
Dihydroquinoline-3-Carboxamide Derivatives
as New Axl Kinase Inhibitors", Journal of
Medicinal Chemistry

Publication

<1 %

30

Park, Young Suk, Jiafu Ji, John Raymond
Zalcberg, Mostafa El-Serafi, Antonio Buzaid,
and Marwan Ghosn. "Oxaliplatin/5-fluorouracil-
based adjuvant chemotherapy as a standard of
care for colon cancer in clinical practice:
Outcomes of the ACCElox registry : Outcomes
of the ACCElox registry", Asia-Pacific Journal
of Clinical Oncology, 2015.

<1 %

-
- 31 Pfirschke, Christina, Christoffer Gebhardt, Inka Zörnig, Maria Pritsch, Stefan B. Eichmüller, Dirk Jäger, Alexander Enk, and Philipp Beckhove. "T cell responses in early-stage melanoma patients occur frequently and are not associated with humoral response", *Cancer Immunology Immunotherapy*, 2015.
Publication
-
- 32 www.ia-cr.ie
Internet Source
-
- 33 Holmberg, L.. "Management of breast cancer in the elderly", *EJC Supplements*, 200403
Publication
-
- 34 Mina, Marco, Renata Boldrini, Arianna Citti, Paolo Romania, Valerio D'Alicandro, Maretta De Ioris, Aurora Castellano, Cesare Furlanello, Franco Locatelli, and Doriana Fruci. "Tumor-infiltrating T lymphocytes improve clinical outcome of therapy-resistant neuroblastoma", *Oncoimmunology*, 2015.
Publication
-
- 35 A. Yasmin, A. Aryadi, A. Widya, A. P. Suwirya et al. " Original ResearchHypertension in rural area: The determinants of left ventricular geometric alteration among hypertensive patients at Kubu Village, Karangasem,

BaliEffectiveness comparison between lisinopril and amlodipine in Melanesian patients with hypertensionAn experimental study of β -D-Glucans extract of polysaccharide peptides as anti-inflammation and anti-oxidant in patients at high-risk of atherosclerosisEchocardiographic findings among hypertensive patients in rural area: The descriptive study at Kubu Village, Karangasem, BaliAddition of left ventricular ejection fraction to the GRACE score in prediction of short term prognosis in patients with non ST-elevation myocardial infarctionEffect of colchicine on HsCRP and mean platelet volume in acute myocardial infarctionCorrelation between GRACE score with spatial QRS-T angle in non-ST elevation myocardial infarction patientsPrevalence of Impaired Kidney Function in Hospitalized Hypertensive Patients in Agoesdjam General Hospital, Ketapang, West Borneo, IndonesiaInappropriate initial treatment in acute coronary syndrome patients: A gap in cardiovascular careCorrelation between plasma histamine level and atherosclerosis severity based on carotid intima media thickness (CMT) measurement in patients with stable coronary artery diseaseClinical characteristics, management and outcome of patients with ST elevation

myocardial infarction in Pekanbaru city:
Preliminary result of single center
experiencePhase II cardiac rehabilitation
program increases functional capacity in post
atrial septal defect surgical closure in adult
patientsThe identification of human telomerase
catalytic subunit in cervical cancer patients and
its potential usage as strategy to reawaken
hibernating myocardium post myocardial
infarctionCould EURO heart risk score predict
the severity of coronary lesion?Pericardial
effusion in systemic lupus
erythematosusIntracardiac metastasis of
hepatocellular carcinomaMyocardial bridging
assessed by computed tomography scan: A
cause of chest pain in patients without
obstructed coronary arteriesCorrelation
between resting heart rate and 6-Minute Walk
Test distance in systolic heart failure
patientsSevere pulmonary hypertension is a
predictor of mortality in pregnant unoperated
atrial septal defectHigh plasma level of soluble
ST2 associated with adverse outcomes during
acute phase of ST-elevation myocardial
infarctionEffect of extracorporeal shockwave
myocardial revascularization on spatial QRS –
T angle resolution in stable coronary artery
disease patientsFragmented QRS study, does
it have any structural abnormality relation?Risk

of coronary artery lesions severity to erectile dysfunction incidence in stable coronary heart disease patients
Clinical profiles and outcomes of patients with ST-Elevation Acute Coronary Syndrome (STE-ACS) in non-primary PCI capable hospital
Correlation between QTc and QTcD with all caused in-hospital mortality after thrombolytic therapy in Dr.Moewardi Hospital, Surakarta
Functional capacity improvement after newly-modified cardiac rehabilitation programe in post-revascularized coronary artery disease patients
Acute heart failure and diabetes are predictors for mortality in 6 months follow up after acute coronary syndrome
Respiratory training as adjuvant of phase 2 exercise program decreased systolic pulmonary artery pressure in post mitral valve surgery patients with residual pulmonary hypertension
Correlation between Tricuspid Regurgitation Duration with Pulmonary Pressure in Heart Failure Patients
Compatibility of myocardial damage observed by echocardiography to MSCT coronary assessment of arterial stenosis in acute coronary syndrome
Hypolipidemic effect and antioxidant activity of (Tamarind) leaves in hypercholesterolemic-fed rats
Correlation between pulmonary hypertension and functional tricuspid regurgitation in Cimacan

General Hospital
Correlation knowledge and attitude regarding behaviour of controlling blood pressure in elderly hypertension patient in Manggar District
Correlation of lipid profile and high sensitivity C-reactive protein among young people
Increased in-hospital mortality associated with hyponatremia in acute heart failure patients
The association between different criteria of obesity to the severity of coronary artery lesion
Predictor postoperative arrhythmias after pediatric cardiac surgery using aortic cross clamp and cardiopulmonary bypass
Profile of hyperthyroid heart disease patients in National Cardiac Center Harapan Kita : A descriptive study
Effect of cardiac rehabilitation programs on functional capacity of post coronary revascularization patient with decrease ejection fraction using Six Minute Walk-Test
Association of postoperative hyperglycemia with in-hospital mortality in patients undergoing on pump coronary artery bypass grafting
A comparison successful reperfusion with and without fibrinolytic therapy in acute ST-segment elevation myocardial infarct patients
Factors influencing ventricular function in repaired (mid-term evaluation)
Correlation between tricuspid regurgitation duration with right ventricular dysfunction in atrial septal defect patients
The

effect of therapy Spiritual Emotional Freedom Technique (SEFT) changes on blood pressure in elderly hypertension experience in Rejosari Karangas Trenggalek 2015Hyponatremia as a predictor of rehospitalization in patients with acute decompensated heart failureEffect of sildenafil therapy to functional capacity in congenital heart disease patient with severe pulmonary hypertension in Sardjito HospitalEisenmenger syndrome is more prevalence in young female patients with uncorrected atrial septal defect: Result of a single-center registryLeuko-glycemic index as an in-hospital prognostic marker in patients with ST-segment elevation myocardial infarction in Haji Adam Malik General HospitalSerum uric acid levels as predictors of incident complications in acute coronary syndromeAnticoagulant therapy in heart failure patients with sinus rhythmEpicardial fat thickness is related to the blood pressure response abnormalities in exercise stress testingRemote ischemic conditioning prior to primary percutaneous coronary intervention: The effect on Six Minutes Walk Test distance and serum NT-ProBNP levelCorrelation between urinary protein and vascular dysfunction before and after Delivery in preeclamptic womenImpact of high hsCRP

level on adverse cardiovascular events and mortality in ST-elevation acute myocardial infarction undergoing fibrinolysis

Correlation between baseline soluble ST2 and global longitudinal strain

2D Speckle tracking echocardiography with left ventricle remodeling post acute myocardial infarction

Difference of peripheral P level between univalvular and multivalvular of rheumatic heart disease

Correlation between the severity of hyperthyroidism with increased of systolic blood pressure in hyperthyroidism patients

Correlation of red cell distribution width with the severity of coronary artery disease in patients with stable coronary artery disease

Chest pain presentation on acute coronary syndrome patients in Dustira Hospital

Result of exercise stress testing after cardiac rehabilitation program after coronary artery bypass grafting surgery

Relationship between plasma level of Endothelin-1 with major adverse cardiac event in ST-elevation myocardial infarction undergoing percutaneous coronary intervention

Controlled clinic blood pressure lowers the risk of left ventricular hypertrophy: An internship study among hypertensive Indonesians in rural areas

Correlation of CHA₂DS -Vasc-HS scores with coronary artery lesions complexity based

on SYNTAX score
The correlation between
body mass index and diastolic indices in non-
diabetic hypertensive male subjects
Risk of
traditional cardiovascular risk factor for the
development of erectile dysfunction in stable
coronary artery disease patients
Relationship
between left ventricle ejection fraction and
submaximal exercise capacity in cardiovascular
patients
Relationship between type 2 diabetes
mellitus with poor heart rate autonomic
response in post acute myocardial infarction
patients
The ECG features among hypertensive
patients in rural area: A descriptive study at
Kubu Village, Karangasem, Bali
Chronotropic
incompetence in coronary artery bypass graft
and heart valve surgery patients: The role of β -
blocker
Correlation between fibrinogen and low
density lipoprotein (LDL) cholesterol levels with
acute myocardial infarction
Cardiac
rehabilitation use among patients with coronary
artery disease on July–December 2015 in Dr.
Hasan Sadikin General Hospital
Bandung
Differences of plasma histamine level
between acute coronary syndrome and stable
coronary artery disease patients
High
Endothelin-1 plasma level gives a tendency
toward increasing rate of in-hospital major
adverse cardiovascular outcomes in ST-
elevation myocardial infarction undergoing

fibrinolysis
Correlation between serum p53 levels and vascular age determined by carotid intima media thickness (CIMT) in patients with intermediate cardiovascular risk factor
Clinical significance of precordial ST segment depression on admission electrocardiogram in patients with acute inferior myocardial infarction
Postprocedural high sensitivity C-reactive protein and the risk of recurrent ischemic events after percutaneous coronary intervention in patients with stable angina pectoris
The messenger ribonucleic acid expression of B-type natriuretic peptide, natriuretic peptide receptor type-A and type-C in cardiomyocytes of obese population
The correlation between serum galectin-3 level with wall motion score index in patients with ST-segment elevation myocardial infarction
Risk factors of atrial fibrillation in patients with chronic heart failure
QT interval prolongation after non-ST elevation myocardial infarction in type 2 diabetic compared with nondiabetic patients
The novel effect of polysaccharide peptides on endothelial dysfunction and dyslipidemia in high risk groups of atherosclerosis
Relation of Left Atrial Spontaneous Echocardiographic Contrast to Neutrophil/Lymphocyte Ratio in Patient With Mitral Stenosis
Analysis of characteristic of

atrial fibrillation versus not atrial fibrillation in
hospitalized heart failure patientsPulmonary
hypertension features in adult atrial septal
defect at RSUP Dr. SardjitoThe association
between plasma tryptase with ventricular
remodeling after 1 month in patients with acute
coronary syndromeThe correlation of plasma
histamine and tryptase with ST-elevation acute
myocardial infarction in acute coronary
syndrome patientsAcute kidney injury after
coronary artery bypass graft surgery: Incidence
and pre operative risk profileThe effect of
diabetes mellitus on in-hospital mortality and
major complications after isolated coronary
artery bypass grafting surgeryFactor those
influence return to work after CABG
revascularization on coronary artery disease
patientsThe correlation of pulmonary vein
endothelin-1 level with pulmonary vascular
resistance before and after mitral valve
surgery with pulmonary hypertensionMitral
valve E-Point septal separation as an
independent predictor for in-hospital mortality
after acute ST-elevation myocardial
infarctionEffects of allopurinol on
complications of post coronary artery bypass
graft (CABG) surgery in coronary artery
disease's patient with left ventricular
dysfunctionMaternal and fetal outcomes of

pregnancy with valvular heart diseaseQRS
duration in ST elevation myocardial infarction
patients undergoing a primary percutaneous
interventionKnowledge of resident medical
officers on hypertensive emergency: A
descriptive studyHyperuricemia and its
association with incident of congestive heart
failure in Madurese populationDiagnostic value
of Duke Treadmill Score in predicting coronary
lesions severity in patients with suspecer
stable coronary artery diseasesRelationship of
plasma glucose levels and troponin-I in patiens
wi...

Publication

36

M. H. Gasem. "Persistence of Salmonellae in
Blood and Bone Marrow: Randomized
Controlled Trial Comparing Ciprofloxacin and
Chloramphenicol Treatments against Enteric
Fever", Antimicrobial Agents and
Chemotherapy, 05/01/2003

<1%

Publication

37

Veenstra, Christine M., Andrew J. Epstein,
Kaijun Liao, Arden M. Morris, Craig E. Pollack,
and Katrina A. Armstrong. "The effect of care
setting in the delivery of high-value colon
cancer care : High-Value Colon Cancer Care",
Cancer, 2014.

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Phyllanthus niruri Linn increase infiltrating lymphocyte and apoptosis of rectosigmoid cancer patients: A Phase II Clinical Trial

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6